

# New prime K-tuple theorem (6)

$$P, P + 4^n (n = 1, \dots, k)$$

Chun-Xuan Jiang

P. O. Box 3924, Beijing 100854, P. R. China

Jiangchunxuan@vip.sohu.com

## Abstract

Using Jiang function we prove that for every positive integer  $k$  there exist infinitely many primes  $P$  such that each of  $P + 4^n$  is prime.

### Theorem

$$P, P + 4^n (n = 1, \dots, k). \quad (1)$$

For every psitive integer  $k$  there exist infinitely many primes  $P$  such that each of  $P + 4^n$  is prime.

**Proof.** We have Jiang function [1]

$$J_2(\omega) = \prod_p [P - 1 - \chi(P)], \quad (2)$$

where  $\omega = \prod_p P$ ,

$\chi(P)$  is the number of solutions of congruence

$$\prod_{n=1}^k [q + 4^n] \equiv 0 \pmod{P}, \quad (3)$$

where  $q = 1, \dots, P - 1$ .

From (3) we have

If  $P < 2k$  then  $\chi(P) = \frac{P-1}{2}$ , if  $2k < P$  then  $\chi(P) = k$ .

Frome (3) and (2) we have

$$J_3(\omega) = \prod_{P=3}^{P < 2k} \frac{P-1}{2} \prod_{2k < P} (P-1-k) \neq 0. \quad (4)$$

We prove that for every positive integer  $k$  there exist infinitely many primes  $p$  such that each of  $P + 4^n$  is prime.

We have the best asymptotic formula [1, 2]

$$\pi_{k+1}(N, 2) = \left| \left\{ P \leq N : P + 4^n = \text{prime} \right\} \right| \sim \frac{J_2(\omega) \omega^k}{\phi^{k+1}(\omega)} \frac{N}{\log^{k+1} N}. \quad (5)$$

where  $\phi(\omega) = \prod_p (P-1)$ .

## References

- [1] Chun-Xuan Jiang, Jiang's function  $J_{n+1}(\omega)$  in prime distribution. (<http://www.wbabin.net/math/xuan2.pdf>)