

Detection of Virtual Photons II

António Saraiva -- 2009-06-28
ajps2@hotmail.com

See Unified Absolute Relativity Theory at:

<http://www.wbabin.net/saraiva/saraiva105.pdf>

<http://www.wbabin.net/saraiva/saraiva223.pdf>

Telepathy is due to virtual photons or longitudinal electromagnetic waves.

$$f = f_0 \sqrt{\frac{2c}{\Delta v}}$$

$$w = \sqrt{c^2 - kf^2} = \sqrt{c^2 - kf_0^2 \frac{2c}{\Delta v}} = \sqrt{c^2 - k_0 f_0^2}$$

$$k_0 = \frac{2ck}{\Delta v} ; \quad \Delta v = k_0 \quad \Leftrightarrow \quad \Delta v = \sqrt{2ck} = 3.4 \times 10^{-13}$$

$$\Delta v = \frac{n+1}{n-1} \frac{kf_0^2}{2c} = 3.4 \times 10^{-13} \quad \Leftrightarrow$$

$$\Leftrightarrow \quad f_0 = 3.44 \times 10^6 \sqrt{\frac{n-1}{n+1}}$$

$$\text{For } f_0 = 5 \times 10^{14} \text{ Hz} \quad \Leftrightarrow \quad n = -1 - 1.4 \times 10^{-8}$$

We need a metamaterial with negative refractive index to detect virtual photons.

The real sixth sense

Why do we have two 3D detectors of acceleration in the head?

Our equilibrium system has two detectors of acceleration, so we can detect gravitational waves.