

## Why Light Is An Ordinary Wave

11/10/2000 Declan Traill ©  
[declan@netspace.net.au](mailto:declan@netspace.net.au)

Light is treated as a different sort of wave than other types of waves, sound waves or water waves for example. This is indicated by the fact that Doppler frequency shift equations used for sound or water waves do not hold for light waves. However, as I shall demonstrate, light waves can be treated as *normal* waves (normal refers to a wave being a disturbance propagating through a medium, rather than a wave existing unsupported in space without a medium) if one takes the Doppler shift equation for light and split it into two component parts.

1. The usual Doppler shift attributed to other types of waves.
2. A relativistic frequency shift of the emitted light at its source.

If the frequency of the emitted light is frequency corrected due to the source's motion prior to applying the normal Doppler shift equation, then the resulting frequency shift is the same as that given by the equation normally used for calculating the Doppler shift for light.

$$\text{Doppler Shift equation for normal waves : } f' = f \times \frac{c}{c + v} \quad (1)$$

$$\text{Doppler Shift equation for light : } f' = f \times \sqrt{\frac{1 - \frac{v}{c}}{1 + \frac{v}{c}}} \quad (2)$$

Where:  $c$  is the transmission speed of the wave.

$v$  is the recession speed of the source.

For a light source moving at speed  $v$ , Relativity states that the frequency of the emitted light will be:

$$f_{emitted} = f \times \sqrt{1 - \frac{v^2}{c^2}} \quad (3)$$

So, treating light as a *normal* wave, the total Doppler shift should be :

$$f' = f_{emitted} \times \frac{c}{c + v} \quad (4)$$

$$f' = f \times \sqrt{1 - \frac{v^2}{c^2}} \times \frac{c}{c+v} \quad (5)$$

To prove that this treatment is valid, it is necessary to show that (2) and (5) are equivalent. This proof follows :

Using (5) gives:

$$\frac{f'}{f} = \sqrt{1 - \frac{v^2}{c^2}} \times \left( \frac{c}{c+v} \right)$$

$$\left( \frac{f'}{f} \right)^2 = \left( 1 - \frac{v^2}{c^2} \right) \times \frac{c^2}{(c+v)^2}$$

$$\left( \frac{f'}{f} \right)^2 = \frac{c^2}{(c+v)^2} - \frac{v^2}{(c+v)^2}$$

$$\left( \frac{f'}{f} \right)^2 = \frac{c^2 - v^2}{(c+v)^2} = \frac{(c+v)(c-v)}{(c+v)^2}$$

$$\left( \frac{f'}{f} \right)^2 = \frac{c-v}{c+v} = \frac{1 - \frac{v}{c}}{1 + \frac{v}{c}}$$

$$\frac{f'}{f} = \sqrt{\frac{1 - \frac{v}{c}}{1 + \frac{v}{c}}}$$

$$f' = f \times \sqrt{\frac{1 - \frac{v}{c}}{1 + \frac{v}{c}}} \quad \text{This is the same as (2)}$$

QED